

Sea quark effects in B_K from $N_f = 2$ clover-improved Wilson fermions

UKQCD Collaboration: J.M. Flynn^{a*}, F. Mescia^{bc†}, A.S.B. Tariq^{ad‡}

^aSchool of Physics & Astronomy, University of Southampton, SO17 1BJ, UK

^bDipartimento di Fisica, Univ. degli Studi “Roma TRE”, via della Vasca Navale 84, I-00146 Roma, Italy

^cINFN, Laboratori Nazionali di Frascati, Via E. Fermi 40, I-00044 Frascati, Italy

^dDepartment of Physics, Rajshahi University, Rajshahi 6205, Bangladesh

We report calculations of B_K using two flavours of dynamical clover-improved Wilson lattice fermions and look for dependence on the dynamical quark mass at fixed lattice spacing. We see some evidence for dynamical quark effects. In particular B_K decreases as the sea quark masses are reduced towards the up/down quark mass. Our meson masses are quite heavy and a firm prediction of B_K is a task for future simulations.

1. INTRODUCTION

B_K determines the value of the $\Delta S = 2$ neutral kaon mixing matrix element according to

$$\langle \bar{K}^0 | Q^{\Delta S=2}(\mu) | K^0 \rangle = \frac{8}{3} B_K(\mu) f_K^2 m_K^2, \quad (1)$$

with μ indicating the scale dependence of the operator $Q^{\Delta S=2}(\mu) = \bar{s}\gamma_\mu(1 - \gamma_5)d \bar{s}\gamma^\mu(1 - \gamma_5)d$.

The quenched lattice value of B_K has more or less settled down (see [1] for a recent review) and the 1997 staggered fermion result [2] $B_K(\overline{\text{MS}}, 2 \text{ GeV}) = 0.63(4)$, corresponding to a renormalisation group invariant (RGI) value $\hat{B}_K = 0.87(6)$, remains the benchmark. The quoted error does not include quenching effects which have been estimated to be as high as $\pm 15\%$ [3] and remain the primary systematic effect to be addressed. Here we report on a calculation [4] using two degenerate flavours of dynamical (clover-improved) Wilson fermions, aiming to look for sea-quark dependence in B_K .

2. SETUP AND SIMULATION

The operator of interest in the continuum is

$$Q_1(\mu) = \bar{s}\gamma_\mu d \bar{s}\gamma^\mu d + \bar{s}\gamma_\mu\gamma_5 d \bar{s}\gamma^\mu\gamma_5 d, \quad (2)$$

*Presenter

†Partially supported by EU IHP-RTN contract HPRN-CT-2002-00311 (EURIDICE)

‡Supported by a Commonwealth Scholarship

which is the parity conserving part of $Q^{\Delta S=2}(\mu)$ in eq. (1). For Wilson fermions, owing to the explicit breaking of chiral symmetry, this operator mixes with other four-fermion operators. Including the overall multiplicative renormalisation, the subtraction of the unwanted operators may be expressed as

$$Q_1(\mu) = Z(\mu, g_0^2) \left(Q_1^{\text{latt}} + \sum_{i \neq 1} \Delta_i(g_0^2) Q_i^{\text{latt}} \right). \quad (3)$$

The coefficients Z and Δ_i have been determined perturbatively to $O(\alpha_s)$ for $\overline{\text{MS}}$ -NDR in [5,6]. The $\Delta S = 2$ lattice operator can now be matched to the continuum one, having the same chiral properties up to $O(a^0)$. There can be remnant operator mixing, beginning at $O(\alpha_s^2 a^0)$, because the Δ_i coefficients are not known exactly. Even though we use an improved-clover action, the four-fermion operators are unimproved and $O(a)$ artefacts can also be present.

By calculating matrix elements with non-zero momentum kaons we introduce another degree of freedom, allowing us partially to remove the lattice artefacts. The chiral behaviour of the matrix element can be written as [3]

$$\langle \bar{P}^0, \mathbf{p} | Q_1(\mu) | P^0, \mathbf{q} \rangle = \alpha' + \beta' m_p^2 + \delta' m_p^4 + p \cdot q \left[\gamma + \gamma' + (\epsilon + \epsilon') m_p^2 + (\xi + \xi') p \cdot q \right] + \dots \quad (4)$$

The parameters γ' , ϵ' and ξ' are corrections to the corresponding physical contributions. In contrast, α' , β' and δ' are artefacts of chiral symmetry breaking, in

Table 1

The configurations used [8,9]. Lattice spacings are fixed from r_0 . m_P/m_V is quoted for $\kappa_{\text{sea}} = \kappa_{\text{val}}$.

β	c_{SW}	κ_{sea}	a/fm	m_P/m_V	#cfs
5.20	2.0171	0.1350	0.103(2)	0.70(1)	100
5.26	1.9497	0.1345	0.104(1)	0.78(1)	100
5.29	1.9192	0.1340	0.102(2)	0.83(1)	80

this case $O(a, \alpha_s^2 a^0)$ with α_s^2 small, which have to be subtracted. In particular the α' term makes B_K diverge in the chiral limit. Following [3], we ignore chiral logarithmic terms in eq. (4), given the errors and large masses used here. For our calculation we neglect higher order terms and use the following expression for the matrix elements:

$$\langle \bar{P}^0, \mathbf{p} | Q_1(\mu) | P^0, \mathbf{q} \rangle = \alpha' + \beta' m_P^2 + (\gamma + \gamma') p \cdot q. \quad (5)$$

We calculate B_K using Clover-improved Wilson fermions [7] on the UKQCD set of unquenched configurations [8,9] listed in table 1. We use three different sea quark masses in the region $m_P/m_V \geq 0.7$ on a volume of $16^3 \times 32$ ($m_P L \geq 7$) with a nearly constant lattice spacing determined from the Sommer scale, r_0 . Propagators and correlators are calculated using the FermiQCD [10,11] code. Five valence quark propagators at $\kappa = 0.1356, 0.1350, 0.1345, 0.1340$ and 0.1335 are generated for each sea quark. We follow the standard procedure [12] to evaluate the matrix element from 3- and 2-point correlation functions.

$$C^{(3)}(t_x, t_y; p_x, p_y; \mu) = \sum_{\mathbf{x}, \mathbf{y}} \langle P_5(\mathbf{x}, t_x) Q(\mathbf{0}; \mu) P_5(\mathbf{y}, t_y) \rangle e^{i\mathbf{p}_x \cdot \mathbf{x}} e^{i\mathbf{p}_y \cdot \mathbf{y}}, \quad (6)$$

$$C_{J_i J_j}^{(2)}(t; p_x) = \sum_{\mathbf{x}} \langle J_i(\mathbf{x}, t) J_j^\dagger(\mathbf{0}, 0) \rangle e^{i\mathbf{p}_x \cdot \mathbf{x}}. \quad (7)$$

We use local pseudoscalar sources J_i for the kaons (smearing with the method of [13] did not improve the signal significantly). In the 3-point functions, Q is at the origin and t_y is fixed at 10 while t_x is varied over the full temporal range. We have checked neighbouring values of t_y but observe no dependence, implying that the ground state is reasonably well isolated. The momentum configurations $\{\mathbf{p}_x, \mathbf{p}_y\}$ are chosen from the pairs $\{(0, 0, 0), (0, 0, 0)\}, \{(0, 0, 0), (1, 0, 0)\}$ and $\{(1, 0, 0), (0, 0, 0)\}$ where an average over equivalent configurations is implied. The matching co-

efficients are evaluated using the boosted coupling, $g_0^2 = 6\langle P \rangle / \beta$, where the average plaquette values are $\langle P \rangle \in \{0.5336, 0.5399, 0.5424\}$ on our ensembles.

3. ANALYSIS AND DISCUSSION

The matrix element is extracted using the ratios:

$$R_3(\mathbf{p}) = \frac{C^{(3)}(t_x, t_y; p_x, p_y; \mu)}{Z_A^2 C_{PP}^{(2)}(t_x; p_x) C_{PP}^{(2)}(t_y; p_y)} \quad (8)$$

$$X(0) = \frac{8}{3} \left| \frac{C_{A_0 P}^{(2)}(t_x)}{C_{PP}^{(2)}(t_x)} \right|^2, \quad X(\mathbf{p}) = X(0) \frac{p_x \cdot p_y}{m_P^2} \quad (9)$$

where Z_A is the axial current renormalisation. Our Method I is to follow [14,15] and fit the equation

$$R_3(\mathbf{p}) = \tilde{\alpha}' + \tilde{\beta}' X(0) + (\tilde{\gamma} + \tilde{\gamma}') X(\mathbf{p}), \quad (10)$$

to find B_K from $\tilde{\gamma}$, neglecting $\tilde{\gamma}'$. This gives an estimate of the leading term in an expansion of B_K for each set of different valence quarks with a given sea quark mass, with the kaons not necessarily being at the physical kaon mass. Using a linear fit versus the unitary pseudoscalar masses, $(am_P)^2|_{\kappa_{\text{sea}}=\kappa_{\text{val}}}$, to go to the up/down limit gives

$$B_K(\overline{\text{MS}}, 2 \text{ GeV}) = 0.49(13), \quad (11)$$

corresponding to the RGI value $\hat{B}_K = 0.69(18)$. In this method the pseudoscalar masses are for valence quarks in the simulated region ($m_q \approx 2m_s$) and the B_K estimate is one where the sea quarks are realistically light but the valence quarks are heavier than the physical strange quark.

In Method II we follow [5,16] and estimate B_K for each $(\kappa_{\text{sea}}, \kappa_{\text{val}})$ combination:

$$\frac{R_3(\mathbf{p}) - R_3(0)}{X(\mathbf{p}) - X(0)} \Big|_{(\kappa_{\text{sea}}, \kappa_{\text{val}})} = B_K(\mu, \kappa_{\text{sea}}, \kappa_{\text{val}}). \quad (12)$$

The resulting values are illustrated in fig. 1. Taking the unitary points (as done in [17]), with $\kappa_{\text{sea}} = \kappa_{\text{val}}$, and extrapolating to the physical kaon mass results in,

$$B_K(\overline{\text{MS}}, 2 \text{ GeV}) = 0.48(13). \quad (13)$$

Here we have $m_P = m_K^{\text{phys}}$, but the sea and valence quarks are degenerate and hence the sea content is not as light as the up/down quarks. Moreover, the kaon comprises two quarks around $m_s/2$.




Figure 1. $B_K(\overline{\text{MS}}, 2 \text{ GeV})$ for each $(\kappa_{\text{sea}}, \kappa_{\text{val}})$ as a function of the squared pseudoscalar mass. Dashed lines are for guidance to distinguish sets with different sea quarks. The filled points joined by a solid line have $\kappa_{\text{sea}} = \kappa_{\text{val}}$. The lightest point (marked by a large cross) is excluded from the analysis.

Although we recognise the presence of lattice artefacts, it seems that dynamical quark effects can lower the value of B_K . Taking this with the observation [18] that $N_f = 2$ results are always below those for $N_f = 0$, a statement also valid for subsequent works, we see that when one has two still-heavy sea quarks, B_K starts to decrease but is consistent with the quenched value. When the sea quarks are taken towards the massless limit, B_K becomes distinctly lower. A similar effect is seen in the domain-wall results of [17,19]. In contrast, preliminary 2+1 flavour results with improved staggered quarks [20] show some sign of a decrease in B_K with decreasing sea-quark mass, but the overall result is consistent with the quenched value. It is intriguing to note that recent Unitarity Triangle fits, which determine $\hat{B}_K$ after imposing the remaining constraints, also find a value below the usual quenched lattice result (for example, $\hat{B}_K = 0.65(10)$ in [21]). However, our primary result is a decrease in B_K as the sea-quark mass decreases, rather than a firm value for use in phenomenology.

Acknowledgment: we thank Ivan Wolton, Ian Hardy, Oz Parchment and the *Iridis* team at the University of Southampton for computing support.

REFERENCES

1. R. Gupta (2003), hep-lat/0303010. Contributed to Workshop on the CKM Unitarity Triangle, CERN, Geneva, Feb 2002.
2. JLQCD Collaboration, S. Aoki et al., Phys. Rev. Lett. 80 (1998) 5271, hep-lat/9710073.
3. S.R. Sharpe, Nucl. Phys. Proc. Suppl. 53 (1997) 181, hep-lat/9609029.
4. UKQCD Collaboration, J.M. Flynn, F. Mescia and A.S.B. Tariq (2004), hep-lat/0406013.
5. R. Gupta, T. Bhattacharya and S.R. Sharpe, Phys. Rev. D55 (1997) 4036, hep-lat/9611023.
6. S. Capitani et al., Nucl. Phys. Proc. Suppl. 63 (1998) 874, hep-lat/9709049.
7. M. Luscher et al., Nucl. Phys. B491 (1997) 323, hep-lat/9609035.
8. UKQCD Collaboration, A.C. Irving, Nucl. Phys. Proc. Suppl. 94 (2001) 242, hep-lat/0010012.
9. UKQCD Collaboration, C.R. Allton et al., Phys. Rev. D65 (2002) 054502, hep-lat/0107021.
10. M. Di Pietro, Nucl. Phys. Proc. Suppl. 106 (2002) 1034, hep-lat/0110116.
11. M. Di Pietro (2000), hep-lat/0011083.
12. M.B. Gavela et al., Nucl. Phys. B306 (1988) 677.
13. UKQCD Collaboration, P. Boyle, J. Comput. Phys. 179 (2002) 349, hep-lat/9903033.
14. SPQCDR Collaboration, D. Becirevic et al., Nucl. Phys. Proc. Suppl. 119 (2003) 359, hep-lat/0209136.
15. M. Crisafulli et al., Phys. Lett. B369 (1996) 325, hep-lat/9509029.
16. R. Gupta et al., Phys. Rev. D47 (1993) 5113, hep-lat/9210018.
17. RBC Collaboration, T. Izubuchi (2003), hep-lat/0310058.
18. A. Soni, Nucl. Phys. Proc. Suppl. 47 (1996) 43, hep-lat/9510036.
19. RBC Collaboration, C. Dawson (2004). These proceedings.
20. HPQCD–UKQCD Collaboration, E. Gamiz et al. (2004), hep-lat/0409049. These proceedings.
21. UTfit Collaboration, M. Bona et al. (2004), hep-ph/0408079.